





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12P 17/16, 17/12, 41/00	A1	(11) International Publication Number: WO 98/58073 (43) International Publication Date: 23 December 1998 (23.12.98)
(21) International Application Number: PCT/US98/11501 (22) International Filing Date: 15 June 1998 (15.06.98) (30) Priority Data: 08/877,675 17 June 1997 (17.06.97) US (71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). (72) Inventors: MORGAN, William, Brian; 31 Rolling Hill Drive, Chatham, NJ 07928 (US). LIU, Jinchu; 11 Palm Court, Edison, NJ 08820 (US). (74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept., K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ENZYMATIC KINETIC RESOLUTION OF AN INTERMEDIATE USEFUL FOR PREPARING SUBSTITUTED TRI-CYCLICS <div style="text-align: center;">  ((+)-I)  ((±)-II) </div> (57) Abstract <p>The invention relates to a process for preparing a substituted (6,11-dihydro-5H-benzo [5,6]cyclohepta[1,2-b]pyridin-11-yl)piperidine compound of formula ((+)-I) wherein: R, R¹, R², R³ and R⁴ are independently selected from the group consisting of hydrogen, halo, C₁-C₆ alkyl, amino, -OCH₃, -OCF₃ and CF₃, and the dotted line represents an optional double bond; comprising: enzymatically catalyzing the acylation of a compound of the formula ((±)-II) wherein the variables are as defined above, and hydrolysing the product to obtain ((+)-I).</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5


10 **ENZYMATIC KINETIC RESOLUTION OF AN INTERMEDIATE
USEFUL FOR PREPARING SUBSTITUTED TRICYCLICS**

BACKGROUND OF THE INVENTION

15 This invention provides an enzymatic process for preparing optically enriched intermediates useful in the preparation of substituted tricyclic compounds known as antihistamines and as inhibitors of farnesyl protein transferase (FPT). In particular, the process of this invention is useful in preparing intermediates useful in the preparation of FPT inhibitors disclosed, for example, in International Publication
20 Number WO95/10516, published April 20, 1995.

The use of enzymes for the synthesis of non-racemic chiral compounds is now well established. Since they are easy to use and readily available, hydrolases (proteases, esterases and lipases) have been used for the preparation of chirally pure molecules, under both
25 aqueous and non-aqueous conditions. Enzyme catalyzed acylation reactions in non-aqueous solvents have been widely used for the kinetic resolution of racemic alcohols and amines. There are numerous examples in the literature of the selective acylation of a single enantiomer of a racemic primary amine:


30 Scheme A:



However, the enzymatic acylation of secondary and cyclic amines has been described less frequently:


-2-

Scheme B:



Most of the examples gleaned from the literature involve the acylation of
 5 chirally pure proline esters or amides catalyzed by alcalase (Chen et al,
Biorg. Med. Chem. Lett., 4 (1994), p. 443), clostridiopeptidase B (Fortier
 et al, *Biotechnol. Lett.*, 8 (1986), p. 777), α -chymotrypsin (Paradkar et al,
J. Amer. Chem. Soc., 116 (1994), p. 5009), and aminoacyl-t-RNA
 synthetase (Nakajima et al, *Int. J. Pept. Protein Res.*, 28 (1986), p. 1986).
 Examples illustrating the enzymatic acylation of chiral secondary amines
 10 are shown in the following reaction schemes:


Scheme C:



(Gutman et al, *Tet. Lett.*, 33 (1992), p. 3943).

15

Scheme D:



PPL Catalyzed Acylation of 2-Hydroxymethylpiperidine


Enzyme	Temp °C/Time h	ees	eep	Conversion	E
PPL	RT/ 4	0.23	0.70	0.25	7
	40/ 4	0.13	0.59	0.18	4
	0-5/ 30	0.39	0.51	0.43	4

(Asensio et al, *Tet. Lett.*, 32 (1991), p. 4197).

20

-3-

Scheme E:




Enzymatic Acylation of 3-Hydroxymethylpiperidine

Solvent	Enzyme mg/mmol	Vinyl Acetate equiv.	Time h	2 (Yield)(ee)	3 (Yield)(ee)
Vinyl Acetate	300	50	91	29% (0.19)	69% (n/d)
Acetonitrile	300	5	94	37% (<0.02)	42% (n/d)
CH ₂ Cl ₂	100	2.5	7.5	68% (0)	-

(Herradon et al, *S. Synlett* (1995), p. 599).

5

Scheme F:



Product	R	% yield	ee
2 a	methyl	24	0.27
2 b	ethyl	31	0.31
2 c	Allyl	49	0.84


Conditions: Substrate, 1 mmol; Carbonate, 1 mL;

Enzyme, 20 mg; RT, 45 h.

(Orsat, et al, *J. Amer. Chem. Soc.*, 118 (1996), p. 712).

10

Scheme G:

15 (Orsat, et al, *J. Amer. Chem. Soc.*, 118 (1996), p. 712).

Generally, the resolutions suffer from low reactivity and/or selectivity.

The reactions shown in schemes D and E probably occur by enzymatic


-4-

acylation of the primary hydroxyl, followed by non-enzymatic intramolecular acyl transfer:

- (In the above reaction Schemes A-H, the designation of the R substituents is for convenience in discussing those reactions, but does not correspond to the designation of the R substituents in the process claimed below.)

SUMMARY OF THE INVENTION


- This invention provides a highly selective process for preparing a substituted (6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]-pyridin-11-yl)piperidine compound of the formula (+)-I



wherein:

- R, R¹, R², R³ and R⁴ are independently selected from the group consisting of hydrogen, halo, C₁-C₆ alkyl, amino, -OCH₃, -OCF₃ and CF₃, and the dotted line represents an optional double bond; comprising:

- (1)(a) enzymatically catalyzing the acylation of a compound of formula (±)-II, wherein the variables are as defined above, to obtain a compound of formula (+)-III



wherein the enzyme is a hydrolase and wherein the acylating agent is of the formula R⁵COOR⁶, wherein R⁵ is C₁-C₁₅ alkyl, halo methyl, aryl, benzyl or benzyloxy, R⁶ is C₁-C₆ alkyl, C₁-C₆ alkenyl, -COR⁷,

-5-

trifluoroethyl, $-\text{CH}_2\text{CH}(\text{OCOR}^7)\text{CH}_2\text{OCOR}^7$, halo methyl or benzyl, and R^7 is $\text{C}_1\text{-C}_{15}$ alkyl; and

(b) hydrolysing the compound of formula (+)-III;

(c) optionally converting an undesired isomer from step (a)

5 wherein a double bond is present to the racemate by heating, and resubjecting the racemate to enzymatic acylation and hydrolysis;

or

(2) enzymatically catalyzing the acylation of a compound of formula (\pm)-IIa, wherein R , R^1 , R^2 , R^3 and R^4 are as defined above and
10 the bond is a single bond, with a hydrolase, and wherein the acylating agent is as defined above.

Preferred compounds of formula (+)-I made by this process are those wherein R^3 is not hydrogen. Also preferred are compounds
15 wherein R is halo. Still another group of preferred compounds is that wherein R^1 is hydrogen and R , R^2 , and R^3 are selected from the group consisting of halo. Halo is preferably Cl or Br.

DETAILED DESCRIPTION

20 As used herein, the term "halo" means fluoro, chloro, bromo and iodo, with chloro and bromo being preferred.

As used herein, the term "aryl" means phenyl, naphthyl, substituted phenyl or substituted naphthyl, wherein "substituted" means
25 1-3 substituents independently selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, halo, NO_2 and halo methyl.

Those skilled in the art recognize that suitable acylating enzymes
• may have opposing selectivity, and therefore may involve either direct or subtractive resolution. That is, some enzymes may acylate the desired isomer, requiring separation of the isomers, followed by hydrolysis to
30 obtain the desired product (i.e., direct resolution, as claimed in step (1)), while others may acylate the undesired isomer, requiring only separation of the isomers (no hydrolysis) to obtain the desired isomer (i.e., subtractive resolution, as claimed in step (2)).

Commercially available enzymes suitable for use in the claimed
35 process include Altus ChiroCLEC™ PC (*Pseudomonas cepacia*);

-6-

Amano Lipase AY-30 (*Candida rugosa*); Meito Lipase MY (*Candida rugosa*), Meito Lipase AL (*Achromobacter sp.*), Meito Lipase QLC (*Alcaligenes sp.*) and Meito Lipase QLG (*Alcaligenes sp.*); Toyobo LIP-300 and LIP-301 (*Pseudomonas sp.*); Novo SP435 and Novozym 435
5 (*Candida antarctica* lipase B); Boehringer Mannheim Lipase (*Pseudomonas sp.*); and Boehringer Chirazyme™ L3 (*Candida rugosa*), Chirazyme™ L4 (*Pseudomonas sp.*) and Chirazyme™ L6 (*Pseudomonas sp.*).

Preferred enzymes are Toyobo LIP-300/301, Altus Chiro CLEC™
10 PC, Boehringer Mannheim Lipase, Novo SP435 and Novozym 435.

Acylating agents of formula R^5COOR^6 are commercially available or can be prepared by known methods. Preferred acylating agents are trifluoroethyl acetate (TFEOAc), trifluoroethyl butyrate (TFEOBu), trifluoroethyl isobutyrate (TFEOiBu), trifluoroethyl benzoate (TFEBenz),
15 triacetin and tributyrin.

The enzymatic acylation may be carried out in a solvent such as an alkyl acetate such as methyl acetate (MeOAc) or isopropyl acetate, t-butyl methyl ether (TBME), tetrahydrofuran (THF), acetone, acetonitrile, t-amyl alcohol, t-butyl alcohol, pyridine or dioxane. Alternatively, the
20 acylating agent may serve as the solvent. A preferred acylating agent which may also act as the solvent is trifluoroethyl isobutyrate.

The reaction is carried out in a temperature range of 0 to 50°C, preferably at 25 to 30°C (e.g., ambient temperature). The reaction time ranges from 18 to 48 hours, with 24 hours being preferred. The enzyme
25 is added at a ratio of about 1: 2 times the amount of the starting material, preferably about 2 times the amount. The acylating agent is present at about 2 to 10 times the starting material, preferably about 3 times the amount of starting material when the enzyme is present at 2 times the amount of the starting material.

30 The hydrolysis is carried out using standard procedures well known in the art. For example, the acylated compound is refluxed with an acid such as H_2SO_4 . The desired isomer is then recovered by precipitating out by adding a base such as NaOH.

The reaction is preferably carried out under anhydrous
35 conditions. The solvent, or acylating agent when used as the solvent,

-7-

can be anhydrous, or the solution of the starting material in the solvent, or acylating agent when used as the solvent, can be dried by azeotropic distillation before the enzyme is added. The enzyme should be dried under vacuum before adding to the solution, preferably to <700 ppm water.


5

The undesired (-) isomer can be recovered from enzymatic isobutyrylation of racemic II. Heating (-)-II in diphenyl ether or diethylene glycol dibutyl ether (5-15:1, v:v) at 200-260°C, preferably 210°C, for 0.5-26 hours results in complete racemization to racemic II which can be recovered in 77-95% yield with 93-99% purity. The recovered racemic II undergoes the enzymatic isobutyrylation under the same conditions as above.


10

Preferred embodiments of the claimed process are shown in the following reaction schemes:


15 Scheme 1:




Scheme 2:



-8-




Scheme 3:




Especially preferred embodiments of the claimed process are
5 represented by the following reaction schemes:

Scheme 1A:



-9-

Scheme 2A:



Previous methods used to resolve isomers to obtain compounds of formula I involved the resolution of a compound of formula (±)-IIa by chiral chromatography or chemical resolution using stoichiometric amounts of a resolving agent. The process claimed herein uses a biocatalyst to effect the resolution, the biocatalyst being reusable up to 15 times. Furthermore, compounds IIb are chiral atropisomers at room temperature due to restricted rotation about the double bond. However, the isomers can be racemized at high temperatures. By carrying out the enzymatic resolution of IIb, the undesired (-) isomer can be isolated, racemized at 200-260°C, preferably at 210°C, and then subjected to a further enzymatic acylation to increase throughput of (±)-IIIa.

The products of this process are intermediates useful in the preparation of tricyclic compounds useful as farnesyl protein transferase inhibitors such as those disclosed in International Publication Number WO95/10516, published April 20, 1995.

The following tables show the results of varying the various parameters of the reactions. In most of the tables, the compound of formula I prepared by the process has the following substituents: R and R³ are each bromo, R¹ and R⁴ are each hydrogen and R² is chloro; those skilled in the art will recognize that compounds with different R-group substitution are expected to react in a similar manner. In the


-10-

tables and elsewhere in this application, the terms have the following meanings: ee_s is the enantiomeric excess of the unreacted starting material; ee_p is the enantiomeric excess of the product; c is the conversion ($ee_s/(ee_s + ee_p)$); E is the Enantiomeric Ratio:

- 5 $(\ln[(1-ee_s)(1-c)]/\ln[(1+ee_s)(1-c)] \text{ or } \ln[1-c(1+ee_p)]/\ln[1-c(1+ee_p)])$;
Ac is acetyl, OAc is acetate, Me is methyl, Et is ethyl, Pr is propyl and TFE is trifluoroethyl.

ENZYMATIC TRANSESTERIFICATION

A. Screen Results



- General Procedure: TFEEOAc (0.06 mL, 20 equivs.) was added to a mixture of (\pm)-IIc (10 mg) and the enzyme (2-100 mg) in TBME (1.0 mL), except for Runs 1 and 4 which were run in the presence of CaCO_3 (30-40 mg) with MeOAc (1.0 mL) as both solvent and acylating agent. The reactions were shaken at 250 rpm at ambient temperature and monitored by thin layer chromatography. Reactions of interest were analyzed by chiral HPLC, the results of which are shown in Table 1.


Table 1. (\pm)-IIc Acetylation Screen: Results from 248 Enzyme Preparations

Run	Enzyme	Wt mg	Time h	ee_s	ee_p	c	E
1	Altus ChiroCLEC PC	4.1	45	0.22	0.60	0.26	5
2	Amano Lipase AY	16	40	0.18	0.19	0.48	2
3	Meito Lipase MY	18	40	0.18	0.17	0.51	2
4	Toyobo LIP-300	7.0	45	0.22	0.66	0.25	6
5	Toyobo LIP-300	18	16	0.93	0.58	0.61	12
6	Boehringer Chirazyme L3	16	40	0.14	0.18	0.43	2
7	Boehringer Chirazyme L4	12	40	0.11	0.20	0.35	2
8	Boehringer Chirazyme L6	15	40	0.03	0.04	0.42	1
9	Meito Lipase AL	29.4	63.5	0.19	0.11	0.64	2
10	Meito Lipase QLC	53.6	63.5	0.55	0.24	0.69	3
11	Meito Lipase QLG	94.2	63.5	0.69	0.34	0.67	4
12	Boehringer <i>Pseudomonas</i> sp.	2	4	0.676	0.687	0.50	11

- 20 Conditions: Runs 2,3, 5-11: (\pm)-IIc, 10 mg; Trifluoroethyl Acetate, 20 equiv; TBME, 1.0 mL; RT, 250 rpm. Runs 1,4: (\pm)-IIc, 10 mg; CaCO_3 , 30-40 mg; MeOAc, 1.0 mL (as solvent and acylating agent); RT, 250 rpm. Run12: (\pm)-IIc, 12 mg; Trifluoroethyl acetate, 5 equiv.; TBME, 1.0 mL; RT, 250 rpm

-11-

B. The Effect of Solvent




General Procedure: TFEOAc (0.2 mL, 40 equivs.) was added to a mixture of (±)-IIc (19-25 mg) and Toyobo LIP-300 (19-25 mg) in the appropriate solvent (2.0 mL). The reactions were shaken at 250 rpm at +4°C and monitored by TLC and chiral HPLC. The results are in Table 2.

Table 2. The Effect of Solvent on the Acetylation of (±)-IIc Using Toyobo LIP-300

Run	Solvent	Time h	ee _s	ee _p	c	E
1	MeOAc	91	0.21	0.46	0.31	3
2	nPrOAc	91	0.04	0.03	0.60	1
3	TBME	91	0.65	0.64	0.50	9
4	Toluene	91	0.01	n/d	n/d	n/d
5	THF	91	0.39	0.51	0.43	4
6	Acetone	91	0.09	0.32	0.21	2
7	MeCN	26	0.24	0.77	0.24	10
		91	0.32	0.66	0.32	7
8	CH ₂ Cl ₂	91	0.01	n/d	n/d	n/d
9	tAmylOH	91	0.03	0.37	0.07	2
10	tBuOH	91	0.04	0.35	0.10	2
11	Pyridine	91	0.02	0.06	0.23	1
12	pDioxane	91	0.13	0.33	0.28	2

Conditions: (±)-IIc, 19-25 mg; Toyobo LIP-300, 19-25 mg; Trifluoroethyl Acetate, 40 equivs; Solvent, 2.0 mL; 250 rpm; +4°C.

C. The Effect of Acylating Agent



General Procedure: The acylating agent (20 equivs.) was added to a mixture of (±)-IIc (20 mg) and enzyme (19-26 mg) in TBME (2.0 mL), except for Run 8 which used MeOAc (2.0 mL) as solvent and acylating agent, Run 11 which used 10 equivalents of acylating agent and Run 18 which used 88 equivalents of acylating agent. The reactions were shaken at 250 rpm at ambient temperature, except Run 11 which was

-12-

shaken at +4°C, and the reactions monitored by TLC and chiral HPLC. With the exception of Runs 4, 5, 10, 11, 17, 18 and 19, all reactions were subjected to workup in which the product and the starting material were separated by preparative TLC and the enantiomeric excesses

5 determined separately. The results are collected in Table 3.

Table 3. The Effect of Acylating Agent on the Acylation of (±)-IIc Using Toyobo LIP-300


Run	Acylating Agent	Time h	ee _s	ee _p	c	E
1	Trifluoroethyl Acetate	23.75	0.92	0.82	0.53	32
		Isolated	0.91	0.84	0.52	35
2	Trifluoroethyl Acetate	15	0.92	0.79	0.54	28
		16.5	0.95	0.80	0.54	33
		Isolated				
3	Trifluoroethyl Acetate	15	0.91	0.82	0.53	31
		16.5	0.93	0.85	0.52	43
		Isolated				
4	Trifluoroethyl Chloroacetate	27.00	0.05	0.47	0.09	3
5	Trifluoroethyl Dichloroacetate	27.00	0.01	0.08	0.08	1
6	Trifluoroethyl Butyrate	27.00	0.87	0.90	0.49	53
		Isolated	0.95	0.77	0.55	28
7	Trifluoroethyl Butyrate	15	0.68	0.88	0.44	33
		17	0.77	0.87	0.47	34
		Isolated				
8	Trifluoroethyl Hexanoate	27.00	0.88	0.80	0.52	26
		Isolated	0.94	0.66	0.59	17
9	Trifluoroethyl Laurate	27.00	0.39	0.83	0.32	16
		Isolated	0.70	0.78	0.47	17
10	Methyl Acetate (neat)	23.75	0.07	0.42	0.14	3
11	Isopropenyl Acetate	7	0.12	0.28	0.31	2
12	Triacetin	23.75	0.37	0.89	0.30	24
		Isolated	0.53	0.93	0.36	48
13	Triacetin	15	0.27	0.84	0.24	14
		20.75	0.38	0.96	0.28	81
		Isolated				
14	Tributyrin	27.00	0.29	0.79	0.27	11
		Isolated	0.55	0.94	0.37	60
15	Tributyrin	15	0.30	0.81	0.27	13
		20.75	0.45	0.84	0.35	18
		Isolated				
16	Trifluoroethyl Benzyl Carbonate	15	0.86	0.71	0.55	16
		20.75	0.90	0.61	0.59	12
		Isolated				
17	Dibenzyl carbonate	15	0.06	n/d	n/d	n/d
18	Ethyl Butyrate	15	0.02	0.59	0.04	4
19	Trifluoroethyl benzoate	26	0.235	0.990	0.192	250

Conditions: (±)-IIc, 20 mg; Toyobo LIP-300, 19-26 mg; TBME, 2 mL; Acylating agent, 20 equivs.; 250 rpm; RT. Exceptions: Run 10, MeOAc as solvent and acylating agent; Run 11, Acylating agent 10 equivs. and +4°C; Run 18, 88 equivs. of acylating agent; Run 19, (±)-IIc, 10 mg; Acylating agent, 5 equivs.

-13-

ENZYMATIC ISOBUTYRYLATION OF (\pm)-IIc

A. Enzyme Survey



General Procedure: A mixture of (\pm)-IIc (25 mg), TFEOiBu (0.04 mL, 5 equivs.), 4Å molecular sieves (25-40 mg) and enzyme (6-27 mg) in TBME (1.0 mL) was shaken at ambient temperature and 250 rpm for 23.5 h. The reactions were monitored by chiral HPLC and the results are collected in Table 4.

Table 4. Isobutyrylation of (\pm)-IIc with Various Enzyme Preparations in TBME

Run	Source	Enzyme	ee _s	ee _p	c	E
1	Sawa	Immobilized Lipase LIP-301 lot# 33580	0.610	0.982	0.383	203
2	Sawa	Lipoprotein Lipase LPL-701 lot#0514A	0.245	0.982	0.200	143
3	Toyobo	Lipase (LIP-300)	0.712	0.985	0.419	278
4	Toyobo	Lipoprotein Lipase (LPL-311) Type A	0.059	0.920	0.060	25
5	Toyobo	Lipoprotein lipase (LPL-701)	0.267	0.979	0.214	123
6	Toyobo	Lipoprotein lipase (Type A)	0.079	0.932	0.078	31
7	Boehringer	chirazyme L4	0.092	0.910	0.092	23
8	Boehringer	chirazyme L6	0.260	0.977	0.210	110
9	Altus	ChiroClec PC	0.053	0.908	0.055	22
10	Toyobo	LIP-300 lot# 36510	0.724	0.987	0.423	335
11	Toyobo	LPL-311 lot# 53250	0.037	0.922	0.038	26

10 Conditions: (\pm)-IIc (25 mg, 50mM), Trifluoroethyl isobutyrate (5 eq), Enzyme (6-27 mg), 4Å Sieves (25-40 mg), TBME (1.0 mL), 250 rpm, RT, 23.5 h.

B. Effect of Solvent

General Procedure: For Runs 1-9, (\pm)-IIc (49-57 mg), 4Å molecular sieves (47-59 mg) and Toyobo LIP-300 (50-55 mg) were suspended in the appropriate solvent (2.0 mL) and trifluoroethyl isobutyrate (0.08 mL, 5 equivs.) added, except for Runs 1-3 where the solvent was used as the acylating agent. The reactions were shaken at 250 rpm at ambient temperature for 22.5 h.

-14-

For Runs 10-25, mixtures of (\pm)-IIc (70 mg), Toyobo LIP-300 (70 mg) and trifluoroethyl isobutyrate (5 equivs.), except Runs 17-23 which used solvent as acylating agent, in the appropriate solvent (2.0 mL) were shaken at 300 rpm and 30°C for 24 h.

5 The results of the chiral HPLC analysis are collected in Table 5.

Table 5. Effect of Solvent on the isoButyrylation of (\pm)-IIc Using Toyobo LIP-300

Run	Solvent	Acylating Agent	ee _s	ee _p	c	E
1	Trifluoroethyl isobutyrate	None	0.445	0.947	0.320	57
2	Ethyl isobutyrate	None	0.106	0.881	0.107	18
3	Methyl isobutyrate	None	0.032	n/d	n/d	n/d
4	TBME	TFEOiBu	0.535	0.984	0.352	217
5	Toluene	TFEOiBu	0.145	0.917	0.137	27
6	THF	TFEOiBu	0.147	0.926	0.137	30
7	Acetone	TFEOiBu	0.097	0.990	0.089	219
8	MeCN	TFEOiBu	0.134	0.924	0.126	29
9	pDioxane	TFEOiBu	0.086	>0.99	0.080	217
10	TBME	TFEOiBu	0.618	0.973	0.388	137
11	10%Et ₃ N/TBME	TFEOiBu	0.851 - 0.917	0.936 - 0.938	0.476 - 0.494	83 - 102
12	10% Toluene/TBME	TFEOiBu	0.541	0.968	0.359	106
13	20% Toluene/TBME	TFEOiBu	0.447	0.967	0.316	93
14	30% Toluene/TBME	TFEOiBu	0.388	0.965	0.287	82
15	40% Toluene/TBME	TFEOiBu	0.316	0.964	0.247	75
16	50% Toluene/TBME	TFEOiBu	0.241	0.966	0.200	72
17	Methyl isobutyrate neat	None	0.022	n/d	n/d	n/d
18	Ethyl isobutyrate neat	None	0.134	0.988	0.119	192
19	10% EtOiBu/TBME	None	0.029	0.956	0.030	46
20	20% EtOiBu/TBME	None	0.059	0.991	0.056	226
21	30% EtOiBu/TBME	None	0.07	0.991	0.066	242
22	40% EtOiBu/TBME	None	0.076	0.991	0.071	251
23	50% EtOiBu/TBME	None	0.092	0.991	0.085	251
24	TBME/Dried Enzyme	TFEOiBu	0.856 - 0.943	0.944 - 0.946	0.476 - 0.499	96 - 130
25	20% Et ₃ N/TBME	TFEOiBu	0.875 - 0.960	0.872 - 0.877	0.501 - 0.523	42 - 60

Conditions: Runs 1-9: (\pm)-IIc (50 mg, 50mM), Toyobo LIP-300 (50-55 mg), TFEOiBu (5 eq., except Runs 1-3 which used solvent as acylating agent), Solvent (2.0 mL), 4Å Sieves (47-59 mg), 250 rpm, RT, 22.25 h.

Runs 10-25: (\pm)-IIc (70 mg, 75 mM), Toyobo LIP-300 (70 mg); Solvent (2.0 mL), TFEOiBu (5 equiv., except Runs 17-23 which used solvent as acylating agent), 30°C, 300 rpm, 24 h.

ENZYMATIC RESOLUTION OF (\pm)-IIId

A. Acylation of (\pm)-IIId Using ChiroCLEC PC




Table 6. Acetylation of (±)-IIId using ChiroCLEC PC in Various Solvents

Run	Solvent	Time h	ee _s	ee _p	Conversion	E
1	EtOAc	20.25	0.02	0	n/d	n/d
2	PrOAc	20.25	0.00	0	n/d	n/d
3	TBME	20.25	0.08	0.67	0.11	6
4	Acetone	20.25	0.07	0.48	0.12	3
5	MeCN	3.75	0.56	0.83	0.40	18
5	MeCN	20.25	0.68	0.78	0.47	17
6	tAmyl Alcohol	20.25	0.02	0.45	0.05	3
7	Pyridine	20.25	0.01	0.20	0.03	2
8	3-Me-3-pentanol	20.25	0.09	1.00	0.08	n/d

5 Conditions: (\pm)-IId (5-9 mg), TFEoAc (12-20 equiv.), CLEC PC (4.5-9.1 mg), Solvent (1.0 mL), RT, 250 rpm.

B. Acylation of (±)-IIId Using Toyobo LIP-300




Table 7. Acetylation of (±)-IIId using Toyobo LIP-300 in Various Solvents


Run	Reaction	Time h	ees	eep	Conversion	E
1	MeOAc	29.75	0.27	>0.95	0.21	>100
2	PrOAc	29.75	0.02	n/d	n/d	n/d
3	TBME	29.75	0.39	0.69	0.36	8
4	Toluene	29.75	0.01	0.27	0.04	2
5	THF	29.75	0.00	n/d	n/d	n/d
6	Acetone	29.75	0.21	0.51	0.29	4
7	MeCN	2.25	0.15	0.82	0.15	12
		4.75	0.31	0.91	0.26	30
		21.75	0.82	0.85	0.49	31
		29.75	0.89	0.80	0.53	26

-16-

8	Dichloromethane	29.75	0.02	n/d	n/d	n/d
9	tAmyl Alcohol	29.75	0.04	0.38	0.11	2
10	Pyridine	29.75	0.02	n/d	n/d	n/d
11	Dioxane	29.75	0.12	n/d	n/d	n/d
12	MeOAc, neat	27.25	0.47	0.92	0.34	41
13	Trifluoroethyl Acetate, neat	27.25	0.17	0.49	0.25	3
14	isoPropenyl Acetate, neat	3	Complete reaction			

Conditions: (±)-IIId, 6.5-12.8 mg; TFEoAc, 0.08 mL, 30-50 equiv.; Enzyme, 2.8-8.4 mg; Solvent, 2.0 mL; RT, 250 rpm., except Runs 12-14 which used solvent (2.0 mL) as acylating agent

C. Isobutyrylation of (±)-IIId Using Toyobo LIP-300




5

Table 8 Isobutyrylation of (±)-IIId Using Toyobo LIP-300 in Various Solvents

Run	Solvent	ees	eep	Conversion	E
1	TBME	0.818	0.971	0.457	174
2	THF	0.474	0.889	0.348	27
3	Toluene	0.202	0.962	0.173	63
4	MeCN	0.236	0.932	0.202	36

Conditions: (±)-IIId, 50 mg; Toyobo LIP-300, 50 mg; TFEoIBu, 0.08 mL, 5 equivs.; Solvent, 2.0 mL; 250 rpm; RT.

D. Acylation of (±)-IIId Using NOVO SP435



10

Table 9. Acetylation of (±)-IIId using Novo SP435 in Various Solvents/Temperatures

Run	Reaction	Time h	ees	eep	Conversion	E
1	MeOAc	15.50	0.45	0.18	0.72	2
2	MeOAc	1.00	0.68	0.70	0.49	11
3	PrOAc	15.5	0.17	0.61	0.22	5
4	TBME	40	0.09	>0.95	0.09	43
5	Toluene	40	0.05	0.85	0.06	13
6	THF	15.5	0.82	0.31	0.72	4
7	Acetone	15.5	0.33	0.78	0.30	11
8	Acetone	1	0.72	0.64	0.53	10
9	MeCN	15.5	0.24	0.02	0.91	1

-17-

10	MeCN	1	>0.95	0.43	0.69	8
11	MeCN, 0°C	1.25	0.55	0.73	0.43	11
12	MeCN/NaHCO ₃ ; -2 to -5°C	1.5	0.39	0.77	0.34	11
13	Dichloromethane	4.0	0.07	>0.95	0.07	42
14	tAmylOH	15.5	0.02	>0.95	0.02	40
15	Pyridine	15.5	0.52	0.35	0.60	3

Conditions: Runs 1,3-7,9, 13-15: (±)-IIId, 7-11 mg; TFEOAc, 50 mL, 14-25 equiv.; SP435, 8-13 mg; Solvent, 2.0 mL, RT, 250 rpm.

Runs 2, 8, 10-12: (±)-IIId, 12-15 mg; TFEOAc, 100 mL, 25-31 equiv.(except Run 2 which used solvent as the acylating agent); SP 435, 4-7 mg; Solvent, 2.0 mL; 250 rpm.

5

Table 10. Acylation of (±)-IIId with Various Acylating Agents catalyzed by Novo SP435

Run	Solvent/ Acylating Agent	Acylating Agent equiv.	Time h	ees	eep	c	E
1	MeOAc	Neat	66.25	>0.95	0.09	0.91	n/d
2	MeOAc	Neat	2.0	0.26	0.83	0.24	13
3	MeOAc	Neat	1.0	0.68	0.70	0.49	11
4	EtOAc	Neat	66.25	>0.95	0.11	0.90	n/d
5	PrOAc	Neat	66.25	0.74	0.47	0.61	6
6	iPropenylOAc (neat)	Neat	66.25	0	0	1.00	n/d
7	iPropenylOAc	12	66.25	0	0	1.00	n/d
8	Acetic Anhydride	10	66.25	0	0	1.00	n/d
9	TFEOAc	14	66.25	0.60	0.29	0.67	3
10	TFEOAc/Acetone -5°C	11	20.75	0.16	0.76	0.18	8
11	EtOAcCl	Neat	66.25	0	0	0	n/d
12	TFEOAcCl	17	66.25	0	0	0	n/d
13	Propionic Anhydride	14	66.25	0	0	1.00	n/d
14	Butyric Anhydride	17	66.25	0	0	1.00	n/d
15	isoButyric Anhydride	17	66.25	0	0	1.00	n/d
16	EtOBu	Neat	66.25	0	0	0	n/d
17	TFEOBu	17	66.25	0.07	0.85	0.07	14
18	TFEOBu/MeCN	12	20.75	0.08	n/d	n/d	n/d
19	TFEOiBu	16	66.25	0	0	0	n/d
20	TFEOiBu/MeCN	11	20.75	0.04	n/d	n/d	n/d
21	MeOAcOMe	53	2.0	0.29	0.40	0.42	3
22	TFEHexanoate	25	5.0	0.29	0.83	0.26	14
23	TFELaurate	15	5.0	0.54	0.88	0.38	26
24	TFE2-MeButyrate	25	5.0	0	0	0	n/d

Conditions: Runs 1, 4-9, 11-17, 19: (±)-IIId, 4.9 mg; TBME or neat acylating agent, 1.0 mL; SP 435, 6.2-10.8 mg; 250 rpm; RT.

Runs 2, 21-24: (±)-IIId, 9-12 mg; MeCN or neat acylating agent, 2.0 mL; SP 435, 5-7 mg; CaCO₃, 42-33 mg, 250 rpm, RT.

Runs 10, 18, 20: (±)-IIId, 11-16 mg; Solvent, 2.0 mL; SP 435, 6-8 mg; NaHCO₃, 34-37 mg (except Run 10), 250 rpm, RT.


Run 3: (±)-IIId, 15 mg; Acylating agent/solvent, 2.0 mL; SP435, 4 mg; 250 rpm; RT

10

15

-18-

E. Other Substrates



Substrate	R ¹	R ²	R ⁴	R ³	ee _s or [α] _D ²⁵	ee _p or [α] _D ²⁵	c	E
1	NH ₂	Cl	H	Br	0.770	0.953	0.447	98
2	H	Me	H	OMe	0.520	0.989	0.345	311
3	H	Cl	NH ₂	Br	-8.09° (c 1.484, MeOH)	+114.1° (c 0.142, MeOH)	n/d	n/d

Conditions:

- 5 Run 1: 1, 5 mg; LIP-301, 10 mg; Trifluoroethyl isobutyrate, 10 equivs.; TBME, 1.0 mL; 200 rpm; RT.
- Run 2: 2, 5.4 mg; LIP-301, 16.6 mg; Trifluoroethyl isobutyrate, 20 equivs.; TBME, 1.0 mL; 200 rpm, RT.
- 10 Run 3: 3, 0.2 g; LIP-301, 0.4 g; Trifluoroethyl isobutyrate, 10 equivs.; TBME, 4 mL; 200 rpm, RT.

Following is a detailed example of a preferred embodiment of the process of this invention.

EXAMPLE 1

- 15 A mixture of (±)-IIc (20 g, 42.7 mmol, 98% pure by assay) in TBME (600 mL) was stirred at ambient temperature for 1 h, then filtered to remove a small amount of insoluble material. The solution was dried by azeotropic distillation; after 200 mL was distilled, a further 200 mL of TBME was added to the reaction mixture. After a total of 400 mL had
- 20 been distilled, the moisture content (Karl-Fischer) of the solution was 214 ppm. Toyobo LIP-300 (40 g; 1282 ppm water) was added to the reaction mixture and stirred for 0.5 h; moisture content at this stage was 250 ppm. Trifluoroethyl isobutyrate (19.1 mL, 3 equivs.) was added and the mixture was stirred at ambient temperature. The reaction was

-19-

terminated after 24 h. The enzyme was removed by filtration and washed with TBME (100 mL).

The combined filtrates were extracted with 0.5M H₂SO₄ (100 mL, 50 mL, 50 mL). The combined acidic extracts were added slowly (60 min) to a mixture of 50% NaOH (15 mL) and water (150 mL). The solid which precipitated was filtered and dried to give (-)-Ic (10.6 g, 96% pure by assay, 52.1% yield; 76.7% ee).


The reaction mixture was extracted with 6M H₂SO₄ (2 X 30 mL). The combined extracts were heated to reflux for 8 h, cooled to room temperature, then added slowly (90 min.) to a mixture of 50% NaOH (70 mL) and ice (170 g), maintaining the temperature at <40°C. The precipitated solid was filtered and dried to give (+)-Ic (8.8 g, 97% pure by assay, 43.4% yield; 98.4% ee).

15

Example 2

(-) II (27.30 g, 94% pure, 81.4% ee) was dissolved in diphenyl ether (137 mL) and heated at reflux under N₂ for 40 min., by which time the ee was <1%.

The mixture was cooled to room temperature and diluted with TBME (500 mL). Analysis of this solution showed a solution yield of 95.8%. The solution was extracted with 0.5 M H₂SO₄ (2 X 218 mL) and the combined acidic extracts were added slowly over a period of 1 hour to a vigorously stirred mixture of 50% NaOH (45 mL) and water (405 mL). After stirring for 0.5 hours, the precipitated solid was collected by filtration and washed with water (820 mL) (26.11 g, 94.6% yield, 1.0% ee).



-20-

Enzymatic Resolution: 1st Cycle

Preparation of the R-isobutyramide (+)-IIIb: (\pm)-IIc (93.0 g, 0.2 mol) was dissolved in TBME (2.0 L) and stirred at room temperature for 1 h. The reaction mixture was filtered, the insoluble material washed with more TBME (~1.0 L), and the volume of filtrate adjusted to 2.9 L. The solution of (\pm)-IIc was then dried by azeotropic distillation, removing 1.0 L of the solvent. The solution was cooled to room temperature and Toyobo LIP-301 (200 g) was added. After stirring at room temperature for 1 h, trifluoroethyl isobutyrate (90 mL, 0.56 mol) was added in one portion.

The reaction was stirred at room temperature under N₂ for 24 h. The enzyme was then removed by filtration and washed with TBME (0.9 L). The combined filtrates were extracted sequentially with three portions of 0.5 M H₂SO₄ (450 mL, 225 mL and 225 mL). These combined acidic extracts contained the unreacted (-)-IIc. The organic layer was then extracted with two portions of 6M H₂SO₄ (135 mL and 135 mL). These combined acidic extracts contained the product isobutyramide (+)-IIIb.

Isolation of (+)-Ic: The combined 6M H₂SO₄ extracts were heated at reflux for 14.5 h, then cooled to room temperature. The reaction mixture was then added slowly to a cold, vigorously stirred mixture of NH₄OH (900 mL) and CH₃CN (270 mL). The solid which precipitated was filtered, washed with water and dried (40.5 g, 43.5%; 0.960 ee).

Isolation of (-)-IIb: The combined 0.5 M H₂SO₄ extracts were added slowly to a cold, vigorously stirred mixture of NH₄OH (450 mL) and CH₃CN (270 mL). The solid which precipitated was filtered, washed with water and dried (40.5 g, 43.5%; 0.966 ee).

Racemization of (-)-IIb. Diphenyl ether (190 mL) was degassed under vacuum for 5-10 min and then purged with N₂ for 5-10 min. (-)-IIb (38 g, 81 mmol) was added and the mixture stirred under N₂ and heated to 245°C. The reaction mixture was maintained at 245°C for 2 h, whereupon racemization was complete. After cooling to room temperature, the reaction mixture was diluted with TBME (570 mL) and filtered. The filtrate was extracted with two portions of 0.5 M H₂SO₄ (190 mL and 95 mL). The extracts were combined, charcoal (19 g) added,

-21-

and the mixture heated to reflux for 1 h. After cooling, the mixture was filtered through Celite and the bed washed with 0.5 M H₂SO₄ (95 mL). The combined filtrates were added slowly to a cold, vigorously stirred mixture of NH₄OH (190 mL) and CH₃CN (114 mL). The solid which
5 precipitated was filtered, washed with water and dried (31.9 g, 84.0%).

A similar procedure can be carried out using diethylene glycol dibutyl ether in place of diphenyl ether, and heating at 210°C for about 12 hours.

Enzymatic Resolution: 2nd Cycle


10 Racemized (±)-IIc (30 g, 64 mmol) was dissolved in TBME (600 mL), filtered and the volume adjusted to 900 mL. The solution was then dried by azeotropic distillation, removing 300 mL solvent. The mixture was cooled and Toyobo LIP-301 (60 g; recovered from 1st cycle above) was added. The mixture was stirred for 1 h, then trifluoroethyl isobutyrate (30
15 mL, 190 mmol) was added. After stirring at room temperature under N₂ for 24 h, the reaction mixture was then filtered and the enzyme cake washed with TBME (300 mL). The combined filtrate was extracted with three portions of 0.5 M H₂SO₄ (150 mL, 75 mL and 75 mL) to remove the unreacted (-)-IIc. The organic layer was then extracted with two
20 portions of 6M H₂SO₄ (45 mL and 45 mL) which were combined and refluxed for 16 h. The cooled reaction mixture was then added slowly to a vigorously stirred, cold mixture of NH₄OH (300 mL) and CH₃CN (90 mL). The precipitated (+)-Ic was filtered, washed with water and dried:
(13 g, 43%; 0.986 ee).

25

-22-

We claim:


1. A process for preparing a substituted (6,11-dihydro-5H-benzo-[5,6]cyclohepta[1,2-b]pyridin-11-yl)piperidine compound of the
 5 formula (+)-I



wherein:

- R, R¹, R², R³ and R⁴ are independently selected from the group consisting of hydrogen, halo, C₁-C₆ alkyl, amino, -OCH₃, -OCF₃ and
 10 CF₃, and the dotted line represents an optional double bond;
 comprising:

(1)(a) enzymatically catalyzing the acylation of a compound of formula (±)-II, wherein the variables are as defined above, to obtain a compound of formula (+)-III



- 15 wherein the enzyme is a hydrolase and wherein the acylating agent is of the formula R⁵COOR⁶, wherein R⁵ is C₁-C₁₅ alkyl, halo methyl, aryl, benzyl or benzyloxy, R⁶ is C₁-C₆ alkyl, C₁-C₆ alkenyl, -COR⁷, trifluoroethyl, -CH₂CH(OCOR⁷)CH₂OCOR⁷, halo methyl or benzyl, and
 20 R⁷ is C₁-C₁₅ alkyl; and

(b) hydrolysing the compound of formula (+)-III;

(c) optionally converting an undesired isomer from step (a)

wherein a double bond is present to the racemate by heating, and resubjecting the racemate to enzymatic acylation and hydrolysis;


25 or

(2) enzymatically catalyzing the acylation of a compound of


-23-

formula (\pm)-IIa, wherein R, R¹, R², R³ and R⁴ are as defined above and the bond is a single bond, with a hydrolase, and wherein the acylating agent is as defined above.

- 5 2. A process of claim 1 wherein the enzyme is Toyobo LIP-300, Toyobo LIP-301, Altus Chiro CLEC™ PC or Novozym 435.
3. A process of claim 1 wherein the acylating agent is selected from the group consisting of trifluoroethyl acetate, trifluoroethyl butyrate, trifluoroethyl isobutyrate, trifluoroethyl benzoate, triacetin and tributyrin.
- 10 4. A process of claim 1 for preparing a compound of formula I wherein R, R² and R³ are halo and R¹ and R⁴ are hydrogen.
- 15 5. A process of claim 1 for preparing a compound of the formula



comprising enzymatically catalyzing the acylation a compound of the formula




- 20 using Toyobo LIP-300, Toyobo LIP-301 or Altus ChiroCLEC™ PC as the enzyme and trifluoroethyl acetate, trifluoroethyl butyrate, trifluoroethyl isobutyrate, trifluoroethyl benzoate, triacetin or tributyrin as the acylating agent, followed by hydrolysis and optionally followed by reconversion of the undesired isomer to the racemate, and resubjecting the racemate to
- 25 enzymatic acylation and hydrolysis.
6. A process of claim 1 wherein in step (c), the undesired isomer from step (a) is converted to the racemate by heating at 200-260°C in diphenyl ether or diethylene glycol dibutyl ether.


-24-

7. A process of claim 5 wherein the undesired isomer is converted to the racemate by heating at 200-260°C in diphenyl ether or diethylene glycol dibutyl ether.

5 8. A process of claim 1 for preparing a compound of the formula




comprising enzymatically catalyzing the acylation a compound of the formula




10 using Toyobo LIP-300, Toyobo LIP-301 or Altus ChiroCLEC™ PC as the enzyme and trifluoroethyl acetate or trifluoroethyl isobutyrate as the acylating agent, followed by hydrolysis.

9. A process of claim 1 for preparing a compound of the formula



15 comprising enzymatically catalyzing the acylation a compound of the formula



20 using Novozyme SP435 as the enzyme and trifluoroethyl acetate, trifluoroethyl butyrate, trifluoroethyl hexanoate, trifluoroethyl laurate or methyl acetate as the acylating agent.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/11501

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12P17/16 C12P17/12 C12P41/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	B. ORSAT ET AL.: "Homocarbonates as Substrates for the Enantioselective Enzymatic Protection of Amines." J. AMER. CHEM. SOC., vol. 118, 1996, pages 712-713, XP002078549 cited in the application see the whole document ---	1-9
Y	B. HERRADON AND S. VALVERDE: "Biocatalysis in Organic Synthesis." SYNLETT, 1995, pages 599-602, XP002078550 cited in the application see the whole document ---	1-9

-/--

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

24 September 1998

Date of mailing of the international search report

07/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Douschan, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/11501

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	G. ASENSIO ET AL.: "Enzyme-mediated Enantioselective Acylation of Secondary Amines in Organic Solvents." TETRAHEDRON LETTERS, vol. 32, no. 33, 1991, pages 4197-4198, XP002000856 cited in the application see the whole document	1-9
Y	--- S.-T. CHEN ET AL.: "Proline as Nucleophile in Kinetically Controlled Peptide Synthesis Catalyzed by Alcalase in 2-Methyl-2-propanol." BIOORG. MED. CHEM. LETT., vol. 4, no. 3, 1994, pages 443-448, XP002054802 cited in the application see the whole document	1-9
Y	--- G. FORTIER AND S. L. MACKENZIE: "Peptide Bond Synthesis by Clostridiopeptidase B." BIOTECHNOLOGY LETTERS, vol. 8, no. 11, 1986, pages 777-782, XP002078551 cited in the application see the whole document -----	1-9